

## OBSERVATIONS ON EYE WORMS OF BIRDS.

By J. W. FIELDING, Australian Institute of Tropical Medicine,  
Townsville, North Queensland.

THE presence of worms in the eyes of birds was apparently first noted about 1819, when Rudolphi described a number of species. From 1825 onwards a noted collector named Natterer appears to have concentrated his attention on the parasites of birds, paying special attention to the parasites of the eyes. He appears to be responsible for quite half the species recorded to date, which were collected principally in Brazil. They were described by Molin (1860), since when various authors have found and described new species from other parts of the world.

Kreff (1871) first recorded the presence of worms from the eyes of wild birds from the Australian region. These worms were placed in the genus *Ascaris* (*Ascaris* sp.), and were recovered from the Red-wattle bird or Gill bird, *Acanthochæra carunculata* Lath. Johnston (1912) redescribed it as *Ceratospira acanthochæra*, later placing it in the genus *Oxyuris*; Railliet, Stiles, and Hassall give the specific name as *Anthochæra*. Von Linstow (1897) found worms in the eyes of Brenchley's Fruit Pigeon, *Zonocnas brenchleyi*, from Bismarck Archipelago, describing it as *Ancyrocanthus ophthalmica*; Ransom (1904) brought it into the genus *Ceratospira*.

From 1913 onward special attention has been paid to the presence of worm parasites in the eyes of wild birds in North Queensland. Nicoll (1914) recorded parasites from the eyes of the Wedge-tail eagle, *Uroæus (Aquila) audax*, and the Brown hawk, *Hieracidea berigora*, which were placed provisionally in the genus *Oxyuris*. Breinl (1913) recorded *Filaria dactylonis* from the Laughing Jackass; *Dacelo leachi*, which was placed by Johnston (1916) in *Ceratospira*; and latterly this species has been placed in the genus *Thelazia* by York and Mapstone (1926). The presence of an *Oxyuris* sp. from the Sea Eagle, *Haliæetus leucogaster* is here recorded. These records form the positive findings of a systematic examination of 250 wild birds for ecto and endo parasites, special attention being paid to the eyes. The list of birds examined includes 109 domestic birds; of the wild birds examined 51 were found to be infected, or 2 per cent. of the total; of the 109 domestic birds 51 were positive, or 46.88 per cent. In the following list of birds examined, the common and ornithological names are given to avoid confusion:—

		Number Exmd.	Nega- tive.	In- fected.
Bee eater (sacred king- fisher)	<i>Halcyon sanctus</i>	1	1	..
Bower bird .. ..	<i>Chlamydodera orientalis</i>	2	2	..
Butcher bird .. ..	<i>Cracticus destructor</i>	3	3	..
Canary, native .. ..	<i>Ptilotis fusca</i>	4	4	..
Cockatoo, black .. ..	<i>Calyptrorhynchus funereus</i>	1	1	..
Cockatoo, white .. ..	<i>Cacatua galerita</i>	6	6	..
Cormorant .. ..	<i>Plotus (Anhinga) novæ hollandiæ</i>	2	2	..
Coot, bald .. ..	<i>Porphyrio melanonotus</i>	19	19	..
Crane, white .. ..	<i>Herodias timoriensis</i>	2	2	..
Crane, blue-grey .. ..	<i>Notaphox novæ hollandiæ</i>	2	2	..
Cuckoo .. ..	<i>Eudynamis cyanocephala</i>	4	4	..
Cuckoo, chestnut breasted	<i>Cacomantis castaneiventris</i>	1	1	..
Cuckoo, shrike, black- faced	<i>Graucalus melanops</i>	2	2	..
Curlew .. ..	<i>Numenius cyanopus</i>	1	1	..
Dove, large (wood-pigeon)	<i>Geopelia humeralis</i>	1	1	..
Drongu, fish tail .. ..	<i>Chibia bracteata</i>	6	6	..
Duck, black .. ..	<i>Anas superciliosa</i>	12	12	..
Duck, Indian Runner (penguin duck)	?	13	13	..
Duck, muscovy .. ..	<i>Cairina moschata</i>	3	..	3
Duck, whistling .. ..	<i>Dendrocygna arcuata</i>	12	12	..
Eagle, sea .. ..	<i>Haliæetus leucogaster</i>	1	..	1
Eagle, wedge tailed .. ..	<i>Uroæus (Aquila) audax</i>	2	1	1
Fig bird .. ..	<i>Sphecotheres maxillaris</i>	11	11	..
Fowl, domestic .. ..	<i>Gallus domesticus</i>	82	34	48
Friar bird (leather head)	<i>Tropidorrhynchus corniculatus</i>	10	10	..
Frog mouth .. ..	<i>Podargus phalacroides</i>	4	4	..

J. Boyd, F.R.G.S.,  
in 1921, have been  
ation of his worth  
nees:—

ity, passed over in  
ralia, after a spell  
he Northern State.  
conducted a school  
and an exceptional  
island Agricultural  
olunteer soldier in  
arrison Artillery.

le entitled, "The  
written by "Nut  
was a Frenchman  
used him of being  
in France at the  
in a stable would  
have been half a  
was prouder than

was largely the  
me, too, of the  
proud. All his  
g, and those in close contact  
Orlando Daly, a tutor at Eton,  
ent Force in Brisbane, and so  
(now Major-General Byron  
of the fact that three of his  
e was in command all became  
Sellheim, General Foott (our  
General Sir Brudenell White—  
ff officers the war produced—  
received some coaching from  
hers of his pupils included  
Postmaster-General), Mr. A.  
ary for Lands), Mr. Horace  
ad others too numerous to

or's writings, he forgot to  
ich he gave an account of  
far-off days that his mind  
These few reminiscences are  
or, schoolmaster, journalist,  
ry activities covered a very  
e was the author of "The  
eology in Verse," a clever  
res on agricultural subjects  
ul outlook on world affairs  
ralian.—Ed. "Q.A.J."]

for fertilising of bananas  
otash as a source of the  
banana culture. Bananas  
rine, and the use of salt  
er muriate of potash for  
d muriate of potash can  
mixtures for bananas.

		Number Exmd.	Nega- tive.	In- fected.
Goose, domestic	<i>Anser cinereus</i>	2	2	..
Goose, pied	<i>Anseranas semipalmata</i>	1	1	..
Grebe, hoary headed	<i>Podiceps poliocephalus</i>	1	1	..
Hawk, brown	<i>Hieracidea berigora</i>	10	9	1
Hawk, sparrow	<i>Accipiter cirrhocephalus</i>	1	1	..
Hen, water	<i>Gallinula tenebrosa</i>	1	1	..
Heron, Nankeen	<i>Nycticorax caledonicus</i>	2	2	..
Honeyeater, blue faced	<i>Entomyza cyanotus</i>	1	1	..
Ibis, glossy	<i>Plegadis falcinellus</i>	3	3	..
Ibis, straw necked	<i>Carphibis spinicollis</i>	2	2	..
Jay, blue	<i>Coracina robusta</i>	4	4	..
Kingfisher, Leach's	<i>Dacelo leachi</i>	10	8	2
Lark, magpie	<i>Grallina picata</i>	7	7	..
Lorikeet, blue-bellied	<i>Trichoglossus novae hollandiae</i>	10	10	..
Magpie	<i>Gymnorhina tibicen</i>	2	2	..
Mocking bird	<i>Anellobia chrysoptera</i>	3	3	..
Mynah bird	<i>Acridotheres tristis</i>	11	11	..
Native companion	<i>Antigone australiana</i>	11	11	..
Oriole, northern	<i>Oriolus chinensis</i>	2	2	..
Owl, brown	<i>Ninox boobook</i>	1	1	..
Owl, masked	<i>Strix novae hollandiae</i>	1	1	..
Parakeet, red shouldered	<i>Neophema pulchella</i>	2	2	..
Pelican	<i>Pelicanus conspicillatus</i>	1	1	..
Pigeon, barred wing	<i>Phaps chalcoptera</i>	2	2	..
Pigeon, domestic	<i>Columba livia domestica</i>	9	7	2
Pigeon, pouter	<i>Macropygia phasianella</i>	4	4	..
Pigeon, purple-crowned	<i>Ptilopus superbus</i>	1	1	..
Plover, spur wing	<i>Lobivanellus lobatus</i>	11	11	..
Rail, land	<i>Eulabeornis philippinensis</i>	2	2	..
Sandpiper	<i>Tringoides hypoleucis</i>	18	18	..
Shag	<i>Phalacrocorax carbo</i>	3	3	..
Sheldrake, white headed	<i>Tadorna radjah</i>	2	2	..
Snipe	<i>Rostratula australis</i>	1	1	..
Spoonbill	<i>Platalea regia</i>	2	2	..
Sun bird	<i>Cinnyris frenata</i>	1	1	..
Swan, black	<i>Cheonopsis atrata</i>	4	4	..
Thrush (babbler)	<i>Pomaterhinus rubeculus</i>	1	1	..
Turkey, scrub	<i>Cathartus lathamii</i>	2	2	..
		359	303	56

#### Examination of Young Chickens.

Attention is drawn to an examination of one poultry yard, taking only young chickens into consideration. The results are not included in the general list of birds examined. The total number examined was forty-five chicks, ranging from three days to twenty-four days old. The table shows the age and the results obtained, giving positive findings in 32 per cent.

	Number Examined.	Negative.	Infected.
3-day old chicks	8	8	0
7-day old chicks	12	12	0
10-day old chicks	14	12	2
14-day old chicks	3	0	3
18-day old chicks	2	0	2
24-day old chicks	8	1	7
10-21-day old muscovy ducks	3	1	2
Total	50	34	16

1 JULY, 1928.] QUEEN

Worm parasites from the at Amoy, China; these were that time this parasite has to be closely associated with following are the places from 1904), Ransom; Jamaica (C of Reunion (Ozoux 1910); B 1916); Hawaii (Norgaard 19 China (Carougeau 1902); 1926), Fielding; New South land (Tryon 1907-8), (Dodd (Fielding 1926); the latter *moschata*.

In a recent paper on the obtained specimens from inf occurs on the sea-coast. Czo tainous districts on Reunion. parasite does not occur on the

As pointed out by Fielding the question of the elucidation on the question for the past two or so experiments and disse tabulated by him, were carried (Leucophaga) *surinamensis* L. and succeeded in infecting young feeding the cockroach to the in the eyes of birds, after having

That there is an association of *parvorum* and the cockroach is evident on consulting the map. This is further accentuated on the cockroach into consideration places where the worm parasite have no records of the occurrence Mauritius, Reunion, New South it has been recorded from and not occur in the places mentioned draw attention to Doucette and portion of the United States, and

#### General Sanitary and Hygienic

It is undoubtedly of first condition of the poultry yard should of poultry keeping the first essential feeding. Infinitely more necessary a disease of such importance as endemic area the poultry yard treating and sterilising effect of use—viz., the sun. This minimum but also as regards shade for The yard should be thoroughly intervals and spread out in the boards, and other material where crickets, and other insect pests at frequent intervals around the or sodium fluoride mixed with prevalent. Recently in the United mended the use of a solution of sodium of the cockroach which is now known worms of poultry. As they were those obtainable in a poultry yard be clearly understood that the bird and all refuse and detritus clean to be a certain amount of risk at scratching around in the soil for with the compound. On general hopper principle, and the drinking

	Number Exmd.	Nega- tive.	In- fected.
ata	2	2	
us	1	1	
	1	1	
	10	9	1
	1	1	
	1	1	
	2	2	
	1	1	
	3	3	
	2	2	
	4	4	
	10	8	2
	7	7	
landice	10	10	
	2	2	
	3	3	
	11	11	
	11	11	
	2	2	
	1	1	
	1	1	
	2	2	
	1	1	
	2	2	
	9	7	2
	4	4	
	1	1	
	11	11	
	2	2	
	18	18	
	3	3	
	2	2	
	1	1	
	2	2	
	1	1	
	4	4	
	1	1	
	2	2	
359	303	56	

poultry yard, taking only young  
added in the general list of birds  
ve chicks, ranging from three  
age and the results obtained,

Number mined.	Negative.	Infected.
8	8	0
2	12	0
4	12	2
3	0	3
2	0	2
3	1	7
3	1	2
	34	16

Worm parasites from the eyes of domestic birds were first found by Dr. Manson at Amoy, China; these were described by Cobbold (1879) as *Flaria mansonii*. Since that time this parasite has been recorded in other parts of the world, and appear to be closely associated with the two tropical lines of Cancer and Capricorn. The following are the places from which the parasite has been recorded:—Florida (Niles 1904), Ransom; Jamaica (Clark 1904), Ransom; Mauritius (Emmerez 1901); Isle of Reunion (Ozoux 1910); Brazil (Megalhaes 1888); Guam Ladrome Islands (Barber 1916); Hawaii (Norgaard 1918); Java (Penning 1894) (Smit 1918); Annam Indo-China (Carougeau 1902); Rabaul Mandated Territory of New Guinea (Heydon 1926), Fielding; New South Wales (Johnston 1909-10); North and Central Queensland (Tryon 1907-8), (Dodd 1909), (Sweet 1910), (Breinl 1913), (Nicoll 1914), and (Fielding 1926); the latter author records finding it in the muscovy duck, *Cairina moschata*.

In a recent paper on the subject, the present writer draws attention to having obtained specimens from inland centres, thereby dispelling the idea that it only occurs on the sea-coast. Ozoux (1910) drew attention to its occurrence in mountainous districts on Reunion. Smit (1918) states that Neveu-Lemaire says that the parasite does not occur on the sea-coast.

As pointed out by Fielding (1926), various experiments have been carried out on the question of the elucidation of the life history, and has himself been working on the question for the past twelve to thirteen years, during which time some thousand or so experiments and dissections, which were the fore-runners of the findings tabulated by him, were carried out. He shows that the cockroach *Pycnocelus* (*Leucophaea*) *surinamensis* L. is responsible for the transmission of the parasite, and succeeded in infecting young and old ducks and young chicks experimentally by feeding the cockroach to the birds, and that the time taken for worms to appear in the eyes of birds, after having swallowed the roaches, is very short.

That there is an association between the eye-worm of poultry *Oxyspirura parvum* and the cockroach *Pycnocelus* (*Leucophaea*) *surinamensis* in Australia is evident on consulting the map showing the distribution so far as is known at present. This is further accentuated on taking the known distribution of *O. mansonii* and the cockroach into consideration throughout the world, which shows that of the places where the worm parasite is known six have also the cockroach. So far we have no records of the occurrence of the roach in Indo-China, Guam, or Rabaul, Mauritius, Reunion, New South Wales, Florida, Jamaica, but owing to the fact that it has been recorded from adjacent places, it is hardly conceivable that it does not occur in the places mentioned. As regards Florida and Jamaica we would draw attention to Doucette and Smith's statement that it occurs on the eastern portion of the United States, and has also been recorded from Mexico.


#### General Sanitary and Hygienic Considerations.

It is undoubtedly of first importance that the general sanitary and hygienic condition of the poultry yard should be attended to. Even under the ordinary rules of poultry keeping the first essentials are—open air, light, cleanliness, and careful feeding. Infinitely more necessary is it that these conditions should be given with a disease of such importance as the one under review. It is considered that in an endemic area the poultry yard should have a minimum of shade so that the penetrating and sterilising effect of nature's own disinfectant can be made of more use—viz., the sun. This minimum of shade is not only meant in the ordinary sense but also as regards shade for insect pests in the fowl-house and yard generally. The yard should be thoroughly cleaned, and the droppings cleaned up at regular intervals and spread out in the sun to dry and eventually burned. All bags, boxes, boards, and other material which might serve as hiding-places for cockroaches, crickets, and other insect pests should be burned. Disinfectants should be sprayed at frequent intervals around the fowl-house and shady spots, or powdered lime, borax, or sodium fluoride mixed with flour should be sprinkled wherever cockroaches are prevalent. Recently in the United States, Doucette and Smith (1926) have recommended the use of a solution of sodium cyanide sprayed on the soil for the destruction of the cockroach which is now known to be responsible for the transmission of eye-worms of poultry. As they were working under conditions totally different from those obtainable in a poultry yard, and as this solution is a deadly poison, it should be clearly understood that the birds would have to be segregated during operations, and all refuse and detritus cleaned up afterwards. Even then there would appear to be a certain amount of risk attached to its use, owing to the fact that the fowls scratching around in the soil for tit-bits may pick up some particles impregnated with the compound. On general grounds the feeding should be carried out on the hopper principle, and the drinking water supplied on the fountain system.



**Treatment.**

Various methods of treatment, aiming at the destruction of the worm parasites of the eyes have been put forward, but to appreciate fully the effects of any treatment, it would appear necessary to point out that there is a wide space all around the eyes communicating with the beak and mouth called the infraocular sinus, through which the tears drop into the buccal cavity, and that the worm can and does pass out from the eyes through this sinus into the mouth, or from one eye to the other, thereby evading for a time the effects of the treatment of the eyes. Obviously, whatever method of treatment is given, it is necessary, to get the best results to carry the treatment to the only means of escape for the worms from the eyes, viz., the opening into the mouth of the sinus. This end is accomplished by painting the opening with the solution by means of a camel-hair brush.



The following solutions have been suggested:—(a) Dropping a solution of sodium bicarbonate into the eyes. (b) A similar proceeding with chloroform. (c) First anaesthetise the eyes with a 5 per cent. solution of cocaine, by drawing apart the eyelids and pouring a few drops of the cocaine into the eyes, allowing this to act for some time, and afterwards drawing up the nictitating membrane and placing a few drops of 5 per cent. creolene underneath. (d) By forcibly extracting the worms by means of a small pair of forceps (this is a dangerous practice and is better left alone). An improvement is here suggested, viz.—a small camel-hair brush. This method has been used for some years and has been found to answer quite well, but requires plenty of time and patience; even this is objectionable

1 JULY, 1928.] QUEENSLAND AGRICULTURAL JOURNAL.

owing to the creation of turpentine and allowing eyes with lukewarm water (Condy's crystals); this for a week or so. Further inflammatory and catarrhal antiseptic as 4 per cent. nine parts of lard and oil

There has been a serious decline in the number of turkeys during recent years, of which is in his annual reports.

This decline in numbers of fresh, as excellent prices, and in seeking the cause of breeding and disease among foxes have probably played a part.

**Suitable Localities.**

The farm, by reason of its location, is well suited to indulge in some of the best turkey raising, and in seeking the cause of breeding and disease among foxes have probably played a part.

**Housing.**

It is natural for a turkey to be in a belt of timber to afford to obtain the best results. The barns are well suited for the purpose, and are protected from the foxes by an area of land, about 15 square feet. The Perches should be placed between the back of the turkey to the number of birds. The Perches should be placed with the help of bushes allowed range during

**Breeding Stock.**

There are several of place. This is a large and appears to be well should only be used for only a year old. One probably six to eight

In selecting, strong from healthy parents and healthy appearance with fair-sized bone

Extra heavy stock is stock in good health thin should never be weakness. Hens will be found to give

owing to the creation of inflammation of the eyes. (e) Placing a small quantity of turpentine and allowing it to act for half an hour, followed by irrigation of the eyes with lukewarm water. (f) A weak solution of potassium permanganate (Condy's crystals); this is somewhat slow in action, and should be continued daily for a week or so. Further treatment directed at the alleviation and cure of the inflammatory and catarrhal conditions by irrigation of the eyes with a mild antiseptic as 4 per cent. boric acid. Ransom suggests the use of a mixture of nine parts of lard and one part of iodoform or carbolsed vaseline.

## TURKEY REARING.

P. RUMBALL, Poultry Expert.

There has been a serious falling off in the number of turkeys kept in Queensland during recent years, of which ample evidence is supplied by the Registrar-General in his annual reports.

This decline in numbers cannot be attributed to the lowered value of turkey flesh, as excellent prices are generally realised for good birds in the local markets, and in seeking the cause, one is forced to the conclusion that faulty methods of breeding and disease are the principal factors, although seasons and the ravages of foxes have probably played their part.

### Suitable Localities.

The farm, by reason of offering turkeys ample range, thereby enabling them to indulge in some of their wild traits, is its natural home. Free range also enables turkeys to gather considerable quantities of their own food in the form of grass, insect life, and when stubbles are available, grains. Districts in which the soil is of a light nature and undulating is more suited to turkey raising than low-lying wet country. Scrub country offers ideal conditions, especially where there is a good supply of green feed and water.

### Housing.

It is natural for turkeys to roost in the open, but, when there is no suitable belt of timber to afford protection, certain housing methods should be adopted to obtain the best results. These houses need not be very elaborate, but should be so constructed as to permit of a free circulation of air. Old open-fronted lofty barns are well suited for this purpose, but in districts in which turkeys have to be protected from the fox it may be advisable to adopt the following system:—Enclose an area of land, about  $\frac{1}{2}$  an acre, with a 6-foot netting fence, and build a shed in the middle. This shed should face north, and be open in front with a 6-inch space between the back of the top wall and roof. The dimensions would vary according to the number of birds to be housed, but each bird should have a floor space of 15 square feet. The house should be 9 feet high in front and 7 feet at back. Perches should be about 3 feet high, all on the same level and 3 inches wide. Suitable nests could be placed around the enclosure and made to look as natural as possible with the help of bushes. The turkeys could be confined to these quarters at night, and allowed range during the day.

### Breeding Stock.

There are several varieties of turkeys, but the American Bronze holds pride of place. This is a large and hardy breed, which has supplanted most other breeds, and appears to be well suited to our climate as well as our markets. Mature stock should only be used for breeders, two years and over being more suited than stock only a year old. One vigorous Tom can be mated with as many as ten hens, but probably six to eight females on the average would give better results.

In selecting, strength and vigour, coupled with the knowledge that your stock are from healthy parents, is of primary importance. The head should have a clean and healthy appearance, body compact and long. Sturdy shanks and strong toes with fair-sized bone indicating stamina.

Extra heavy show specimens do not make the best breeders. What is required, is stock in good hard condition and not fat; on the other hand, stock that are thin should never be used, as the lack of condition may be due to some inherited weakness. Hens weighing 16 to 18 lb. and Toms 25 to 30 lb. in fair condition will be found to give satisfactory results.

OCEAN  
PACIFIC  
SOUTH  
Brisbane  
Cleveland  
Rockhampton  
Tyrone

VOLUME XXX



# QUEENSLAND AGRICULTURAL JOURNAL

Issued by direction of

The Hon. the Secretary for Agriculture

THE UNIVERSITY OF  
QUEENSLAND

Edited by J. F. F. REID

TOWNSVILLE  
COLLEGE  
LIBRARY

JULY to DECEMBER, 1928